

Binod Kumar Sahoo is presently with NIT-  
Rourkela India  
This paper is Archived in Dspace@nitr  
<http://dspace.nitrkl.ac.in/dspace>

Journal of Combinatorial Theory, Series A  
Volume 114, Issue 1 (2007), 52–64

## A characterization of finite symplectic polar spaces of odd prime order

Binod Kumar Sahoo<sup>1</sup>  
and  
N. S. Narasimha Sastry

Statistics and Mathematics Unit  
Indian Statistical Institute  
8th Mile, Mysore Road  
R. V. College Post  
Bangalore-560059, India

E-mails: binodkumar@gmail.com, nsastry@isibang.ac.in

**Abstract.** A sufficient condition for the representation group for a nonabelian representation (Definition 1.1) of a finite partial linear space to be a finite  $p$ -group is given (Theorem 2.9). We characterize finite symplectic polar spaces of rank  $r$  at least two and of odd prime order  $p$  as the only finite polar spaces of rank at least two and of prime order admitting nonabelian representations. The representation group of such a polar space is an extraspecial  $p$ -group of order  $p^{1+2r}$  and of exponent  $p$  (Theorems 1.5 and 1.6).

---

<sup>1</sup>Supported by DAE grant 39/3/2000-R&D-II (NBHM fellowship), Govt. of India

## 1. Introduction

A *point-line geometry* is a pair  $S = (P, L)$  consisting of a nonempty ‘point-set’  $P$  and a nonempty ‘line-set’  $L$  of subsets of  $P$  of size at least 2.  $S$  is a *partial linear space* if any two distinct points  $x$  and  $y$  are contained in at most one line. Such a line, if it exists, is written as  $xy$ ,  $x$  and  $y$  are said to be *collinear* and written as  $x \sim y$ . If  $x$  and  $y$  are not collinear we write  $x \not\sim y$ . The graph with vertex set  $P$ , two distinct points being *adjacent* if they are collinear in  $S$ , is the *collinearity graph*  $\Gamma(P)$  of  $S$ . We write  $d(x, y)$  to denote the distance between two vertices  $x$  and  $y$  in  $\Gamma(P)$ . For  $x \in P$  and  $A \subseteq P$ , we define  $x^\perp = \{x\} \cup \{y \in P : x \sim y\}$  and  $A^\perp = \bigcap_{x \in A} x^\perp$ .  $S$  is *nondegenerate* if  $P^\perp$  is empty. A subset of  $P$  is a *subspace* of  $S$  if any line containing at least two of its points is contained in it. The empty set, singletons, the lines and  $P$  are all subspaces of  $S$ . For a subset  $X$  of  $P$  the *subspace*  $\langle X \rangle$  generated by  $X$  is the intersection of all subspaces of  $S$  containing  $X$ . A subspace is *singular* if each pair of its distinct points is collinear. A *geometric hyperplane* of  $S$  is a subspace of  $S$  different from  $P$ , that meets every line nontrivially.

**1.1. Representations of partial linear spaces.** Let  $p$  be a prime. Let  $S = (P, L)$  be a partial linear space of order  $p$ , that is, each line has  $p+1$  points. (Note that, usually, order of a generalized polygon means something else, see [20], Section 1.3, p. 387).

**Definition 1.1.** (Ivanov [12], p. 305) *A representation of  $S$  is a pair  $(R, \psi)$ , where  $R$  is a group and  $\psi$  is a mapping from the set of points of  $S$  into the set of subgroups of order  $p$  in  $R$ , such that the following hold:*

- (i)  $R$  is generated by the subgroups  $\psi(x), x \in P$ .
- (ii) For each line  $l \in L$ , the subgroups  $\psi(x), x \in l$ , are pairwise distinct and generate an elementary abelian  $p$ -subgroup of order  $p^2$ .

The group  $R$  is then called the *representation group*. The representation  $(R, \psi)$  is *faithful* if  $\psi$  is injective. For each  $x \in P$ , we fix a generator  $r_x$  of  $\psi(x)$  and denote by  $R_\psi$  the union of the subgroups  $\langle r_x \rangle, x \in P$ . A representation  $(R, \psi)$  of  $S$  is *abelian* or *nonabelian* according as  $R$  is abelian or not. Unlike here, ‘nonabelian representation’ in [12] means that ‘the representation group is not necessarily abelian’. A representation  $(R_1, \psi_1)$  of  $S$  is a *cover* of the representation  $(R_2, \psi_2)$  of  $S$  if there exist an automorphism  $\beta$  of  $S$  and a group homomorphism  $\varphi : R_1 \rightarrow R_2$  such that  $\psi_2(\beta(x)) = \varphi(\psi_1(x))$  for every  $x \in P$ . Further, if  $\varphi$  is an isomorphism then the two representations  $(R_1, \psi_1)$  and  $(R_2, \psi_2)$  are *equivalent*.

We now indicate various possibilities for the representation group. Embeddings of partial linear spaces (like projective spaces, polar spaces, generalized polygons, etc.) of order  $p$  in projective spaces over the field  $F_p$  of order  $p$  are all examples of abelian representations. The representation group is the corresponding vector space considered as an abelian group. Every representation of a projective space is faithful (by Definition 1.1(ii)) and the representation group of a finite

projective space of dimension  $m$  over  $F_p$  is an elementary abelian group of order  $p^{m+1}$ . However, a representation of a generalized quadrangle need not be faithful. For example, let  $S = (P, L)$  be a  $(2, 1)$ -generalized quadrangle, let  $P_1, P_2, P_3$  be three triads partitioning  $P$  and let  $R = \{1, r_1, r_2, r_3\}$  be the Klein four group. Define  $\psi : P \rightarrow R$  by  $\psi(x) = \langle r_i \rangle$  if  $x \in P_i$ . Then  $(R, \psi)$  is an abelian representation which is not faithful.

Root group geometries are some examples of nonabelian representations of partial linear spaces. Let  $H$  be a finite simple group of Lie type defined over  $F_p$ . Let  $\mathcal{G} = (P, L)$  be the root group geometry of  $H$ . That is, the ‘point set’  $P$  is the collection of all (long) root subgroups of  $H$ . Two distinct root subgroups  $x, y \in P$  are *collinear* if they generate an elementary abelian subgroup of order  $p^2$  and each subgroup of order  $p$  in it is a member of  $P$ . The ‘line’  $xy$  is the set of  $p + 1$  subgroups of order  $p$  in  $\langle x, y \rangle$ . The identity map defines a representation of  $\mathcal{G}$  in  $H$  and  $H$  is a representation group of  $\mathcal{G}$ . Note that if  $H$  is of type  $E_6, E_7$  or  $E_8$ , then  $\mathcal{G}$  is a parapolar space (see [4], p. 75); if it is of type  $G_2$  or  ${}^3D_4$ , then  $\mathcal{G}$  is a generalized hexagon with parameters  $(p, p)$  and  $(p, p^3)$  respectively (see ([6], p. 322 and 328) for  $p$  odd and ([7], Lemma 2.2, p. 2) for  $p = 2$ ); if it is type  $F_4$  or  ${}^2E_6$ , then  $\mathcal{G}$  is a metasymplectic space (see Section 4, [6]); and if it is of type  ${}^2F_4$ , then  $\mathcal{G}$  is a  $(2, 8)$ -generalized octagon (see [19]). For a discussion of root group geometries including the classical ones, see [5] and [10], Chapter 4.

The following example shows that the representation group for a nonabelian representation of a finite partial linear space could be infinite.

**Example 1.2.** *Let  $S = (P, L)$  be a  $(2, 2)$ -generalized hexagon. Then  $S$  is isomorphic to  $H(2)$  (the one admitting an embedding in  $O_7(2)$ ) or its dual  $H(2)^*$  (see [20], Theorem 4, p. 402). For each  $x \in P$ ,  $H(x) = \{y \in P : d(x, y) < 3\}$  is a geometric hyperplane of  $S$ . The subgraph of  $\Gamma(P)$  induced on the complement of  $H(x)$  in  $P$  is connected if  $S \simeq H(2)$  and has two components if  $S \simeq H(2)^*$  (see [9], section 3). By ([12], Lemma 3.6, p. 310),  $H(2)^*$  admits a nonabelian representation whose representation group is infinite. In fact, this representation is the cover of all other representations of  $H(2)^*$ .*

Our basic tool in this paper (Theorem 2.9) in fact is a sufficient condition on  $S$  and on the nonabelian representation of  $S$  to ensure that the representation group is a finite  $p$ -group.

We refrain from listing several natural questions that suggest themselves regarding the representations and the possible representation groups of finite partial linear spaces. For more on nonabelian representations, see [12].

**1.2. Polar spaces.** A *polar space* [2] here is a nondegenerate point-line geometry  $S = (P, L)$  with at least three points per line satisfying the ‘one or all’ axiom:

*For each point-line pair  $(x, l)$ ,  $x \notin l$ ,  $x$  is collinear with one or all points of  $l$ .*

(see [2], Theorem 4, p. 161 and [22], 7.1, p. 102). *Rank* of  $S$  is the supremum of the lengths  $m$  of chains  $Q_0 \subsetneq Q_1 \subsetneq \cdots \subsetneq Q_m$  of singular subspaces in  $S$ . Since  $L$  is nonempty, the rank of  $S$  is at

least two, but could be infinite. A remarkable discovery of Buekenhout and Shult is that a polar space is a partial linear space ([2], Theorem 3, p. 161). A polar space of rank 2 is a *generalized quadrangle* (GQ, for short). That is, it is a nondegenerate partial linear space such that:

Whenever  $x \in P, l \in L$  with  $x \notin l$ ,  $x$  is collinear with exactly one point of  $l$ .

If a finite GQ has a line with at least three points and a point on at least three lines then there exist integers  $s$  and  $t$  such that each line contains  $s + 1$  points and each point is on  $t + 1$  lines ([3], Theorem 7.1, p. 98). In that case we say that it is a  $(s, t)$ -GQ.

Building on the work of Veldkamp, Tits classified polar spaces whose rank is finite and at least three [22]. (For polar spaces of possibly infinite rank, see [14].) This implies that a finite polar space of rank  $r \geq 3$  and of order  $p$  is isomorphic to either the symplectic polar space  $W_{2r}(p)$  or one of the orthogonal polar spaces  $Q_{2r}^+(p)$ ,  $Q_{2r+1}(p)$  and  $Q_{2r+2}^-(p)$ . For notation see ([21], p. 329). If  $r = 2$  the above yield  $(p, p)$ -,  $(p, 1)$ -,  $(p, p)$ - and  $(p, p^2)$ -GQs respectively. We note the number of points of these polar spaces ([21], Theorem 1, p. 330):

$$\begin{aligned} |W_{2r}(p)| &= (p^{2r} - 1)/(p - 1); \\ |Q_{2r}^+(p)| &= (p^{r-1} + 1)(p^r - 1)/(p - 1); \\ |Q_{2r+1}(p)| &= (p^{2r} - 1)/(p - 1); \\ |Q_{2r+2}^-(p)| &= (p^r - 1)(p^{r+1} + 1)/(p - 1). \end{aligned}$$

The following inductive property of these spaces is important for us (see [3], section 6.4, p. 90).

**Lemma 1.3.** *Let  $S$  be one of the above polar spaces of finite rank  $r \geq 3$  and let  $x, y$  be two noncollinear points. Then  $\{x, y\}^\perp$  is a polar space of rank  $r - 1$  and is of the same type as  $S$ .*

Finite GQs are classified only for  $s = 2, 3$  (see [20], 5.1, p. 401). See [16] for several examples of finite GQs. In [15], Kantor studied finite  $(p, t)$ -GQs  $S$  with  $t \geq 2$  admitting a rank 3 automorphism group  $G$  on points and proved that one of the following holds: (i)  $t = p^2 - p - 1$  and  $p^3 \nmid |G|$ ; (ii)  $G \cong PSp(4, p)$  or  $PFU(4, p)$  and  $S$  is one of the natural GQs associated with these groups; (iii)  $p = 2$ ,  $G = Alt(6)$  and  $S$  is the GQ associated with  $PSp(4, 2)$  ([15], Theorem 1.1). This paper started with a search for new finite  $(p, t)$ -GQs embedded in groups and resulted in a characterization of finite symplectic polar spaces  $W_{2r}(p)$  of rank  $r \geq 2$  for odd primes  $p$  (Theorems 1.5 and 1.6).

**1.3. Extraspecial  $p$ -groups and Hall-commutator formula.** A finite  $p$ -group  $G$  is *extraspecial* if its Frattini subgroup  $\Phi(G)$ , the commutator subgroup  $G'$  and the center  $Z(G)$  coincide and have order  $p$ . An extraspecial  $p$ -group is of order  $p^{1+2m}$  for some integer  $m \geq 1$ , has exponent at most  $p^2$  if  $p$  is odd and 4 if  $p = 2$ , and the maximum of the orders of its abelian subgroups is  $p^{m+1}$  (see [8], section 20, p. 78,79). We denote by  $p_+^{1+2m}$  an extraspecial  $p$ -group of order  $p^{1+2m}$  if its exponent is  $p$  when  $p$  is odd and the abelian subgroups of order  $p^{m+1}$  are elementary abelian when  $p = 2$ . Note that  $p_+^{1+2}$  is isomorphic to the group of  $3 \times 3$  upper triangular matrices with entries from  $F_p$  and 1

on the diagonal. For more on extraspecial  $p$ -groups, see ([11], section 3, p. 127 and Appendix 1, p. 141).

For elements  $g_1, g_2$  in a group, we write  $[g_1, g_2] = g_1^{-1}g_2^{-1}g_1g_2$  and  $g_1^{g_2} = g_2^{-1}g_1g_2$ . We repeatedly use the following Hall's commutator formula ([8], 7.2, p. 22), mostly without mention.

**Lemma 1.4.** *Let  $G$  be a group. Then for  $g_1, g_2, g_3 \in G$ ,*

- (i)  $[g_1g_2, g_3] = [g_1, g_3]^{g_2}[g_2, g_3]$ ;
- (ii)  $[g_1, g_2g_3] = [g_1, g_3][g_1, g_2]^{g_3}$ .

**1.4. Statement of main results.** In this paper we prove:

**Theorem 1.5.** *Let  $S = (P, L)$  be a finite polar space of rank  $r \geq 2$  and of prime order  $p$ . If  $S$  admits a nonabelian representation  $(R, \psi)$  then:*

- (i)  $p$  is odd;
- (ii)  $R = p_+^{1+2r}$ ;
- (iii)  $S$  is isomorphic to  $W_{2r}(p)$ .

**Theorem 1.6.**  $W_{2r}(p)$ ,  $r \geq 2$ , admits a nonabelian representation. Any two such representations are equivalent.

In Section 2 we prove a sufficient condition for a nonabelian representation group to be a  $p$ -group (Theorem 2.9) which is crucial here and also in [18]. In Section 3 we prove Theorem 1.5(i) and that  $R \simeq p_+^{1+2m}$  for some  $m \geq 1$ . In Section 4 we prove Theorem 1.5 when the rank is two. Finally, in Section 5 we prove Theorem 1.5 for the general rank and Theorem 1.6.

## 2. Initial Results

Let  $S = (P, L)$  be a partial linear space. We assume that  $\Gamma(P)$  is connected and that with each  $x \in P$  is associated a geometric hyperplane  $H(x)$  in  $S$  containing  $x$ . Consider the following conditions on  $S$ :

- (C1) If  $y \in H(x)$  then  $x \in H(y)$ .
- (C2) The subgraph  $\Gamma(H'(x))$  of  $\Gamma(P)$  induced on the complement  $H'(x)$  of  $H(x)$  in  $P$  is connected.
- (C3) If  $y \in H'(x)$  then there exist lines  $l_1$  and  $l_2$  containing  $x$  and  $y$  respectively such that for each  $w \in l_1$ ,  $H(w)$  intersects  $l_2$  at exactly one point. Further, this correspondence is a bijection from  $l_1$  to  $l_2$ .
- (C4) The graph  $\Sigma(P)$  with vertex set  $P$  in which two points  $x$  and  $y$  are adjacent if  $y \in H'(x)$  is connected.

**Example 2.1.** *Let  $S = (P, L)$  be a polar space of rank  $r \geq 2$ . Then  $\Gamma(P)$  is connected. For each  $x \in P$ , associate the geometric hyperplane  $x^\perp$  of  $S$ . Then (C1),  $\dots$ , (C4) hold.*

**Example 2.2.** Let  $S = (P, L)$  be a near  $2n$ -gon,  $n \geq 2$ , admitting quads (see [1]). We assume that each line of  $S$  contains at least three points. By definition,  $\Gamma(P)$  is connected. For each  $x \in P$ , associate the geometric hyperplane  $H(x) = \{y \in P : d(x, y) < n\}$  of  $S$ . Clearly (C1) holds. The second corollary to ([1], Theorem 3, p. 155) implies that (C2) holds. Now, ([1], Theorem 2, p. 151) implies that if  $d(x, y) = n$ ,  $x, y \in P$  and  $l_1$  is any line containing  $x$ , then there exists a line  $l_2$  containing  $y$  such that (C3) holds. This also implies that if  $u \sim v$ ,  $u, v \in P$ , then there exists  $w \in P$  such that  $d(u, w) = d(v, w) = n$ . So  $u, w, v$  is a path in  $\Sigma(P)$ . Then connectedness of  $\Sigma(P)$  follows from that of  $\Gamma(P)$ . Thus C(4) holds.

We study nonabelian representations of finite polar spaces of order  $p$  here (Theorems 1.5 and 1.6) and that of near hexagons of order two and admitting quads in [18].

**Remark 2.3.** If  $S = (P, L)$  is a generalized  $2n$ -gon and  $H(x), x \in P$ , is as in Example 2.2, then (C2) need not hold, see Example 1.2.

Let  $(R, \psi)$  be a representation of  $S$ . For  $x, y \in P$ , define  $u_{xy} = [r_x, r_y]$ . Throughout this section we assume that

$$u_{xy} = 1 \text{ whenever } x \in P \text{ and } y \in H(x).$$

**Proposition 2.4.** Assume that (C1) and (C2) hold in  $S$ . Then the following hold:

- (i) If  $u_{vw} = 1$  for  $v, w \in P$  with  $v \in H'(w)$ , then  $r_w \in Z(R)$ .
- (ii) If  $a \in P$  and  $r_a \in Z(R)$ , then  $r_c \in Z(R)$  for every  $c \sim a$ .

*Proof.* (i) Let  $y \in H'(w)$ ,  $y \sim v$  and  $vy \cap H(w) = \{x\}$ . Then  $u_{wy} = 1$  because  $x \notin \{v, y\}$  and  $u_{wx} = u_{vw} = 1$ . Now, connectedness of  $\Gamma(H'(w))$  implies that  $u_{wz} = 1$  for every  $z \in H'(w)$ . Since  $u_{wz} = 1$  for  $z \in H(w)$  also,  $r_w \in Z(R)$ .

(ii) By definition,  $H(a) \subsetneq P$ . Let  $b \in H'(a)$ . By (C1),  $a \in H'(b)$ . By (i),  $r_b \in Z(R)$  because  $u_{ab} = 1$ . Now,  $ac \cap H(b)$  is a singleton. Since each line contains at least 3 points, there exists a point  $z$  in  $ac \cap H'(b)$  different from  $a$ . Now,  $b \in H'(z)$  by (C1) and  $u_{bz} = 1$ . So,  $r_z \in Z(R)$  by (i) again. So the subgroup generated by  $\psi(ac)$  is contained in  $Z(R)$  and  $r_c \in Z(R)$ .  $\square$

**Corollary 2.5.** Assume that (C1) and (C2) hold in  $S$ . If  $R$  is nonabelian then the following hold:

- (i)  $u_{xy} \neq 1$  whenever  $x, y \in P$  and  $y \in H'(x)$ .
- (ii)  $R_\psi \cap Z(R) = \{1\}$ .
- (iii) If  $x \sim y$  then  $y \in H(x)$ .
- (iv) If  $H(x) \neq H(y)$  for each pair of noncollinear points  $x$  and  $y$ , then  $\psi$  is faithful.

*Proof.* (i) follows from Proposition 2.4 and the connectedness of  $\Gamma(P)$ . (ii) and (iii) follow from (i). We now prove (iv). Suppose that  $\langle r_x \rangle = \langle r_y \rangle$  for distinct  $x, y$  in  $P$ . Then  $x \approx y$  by Definition 1.1(ii). By (i),  $u \in H(x)$  if and only if  $u \in H(y)$ . So  $H(x) = H(y)$ , a contradiction.  $\square$

**Proposition 2.6.** *Assume that (C3) holds in  $S$ . Then for  $x, y \in P$ ,  $[u_{xy}, r_x] = [u_{xy}, r_y] = 1$ . If  $u_{xy} \neq 1$  then  $u_{xy}$  is of order  $p$  and  $\langle r_x, r_y \rangle = p_+^{1+2}$ .*

*Proof.* Let  $x \in P$ ,  $y \in H'(x)$  and  $l_1, l_2$  be lines as in (C3). Let  $x, a, u$  be three pairwise distinct points in  $l_1$  and  $y, b, v$  be points in  $l_2$  such that  $y \in H(a)$ ,  $b \in H(x)$  and  $v \in H(u)$ . By (C3),  $y, b, v$  are pairwise distinct. Write  $r_x = r_a^i r_u^j$ ,  $r_y = r_v^k r_b^m$  for some  $i, j, k, m$ , ( $1 \leq i, j, k, m \leq p-1$ ). Now,

$$u_{xy} = [r_a^i r_u^j, r_y] = [r_u^j, r_y] = [r_u^j, r_v^k r_b^m] = [r_u^j, r_b^m] = [r_x r_a^{-i}, r_b^m] = [r_a^{-i}, r_b^m].$$

Since  $[r_a^{-i}, r_b^m] = [r_b^m, r_a^i] r_a^{-i}$ ,

$$\begin{aligned} u_{xy} &= [r_b^m, r_a^i] r_a^{-i} = [r_y r_v^{-k}, r_a^i] r_a^{-i} = [r_v^{-k}, r_a^i] r_a^{-i} = [r_v^{-k}, r_u^{-j} r_x] r_a^{-i} \\ &= [r_v^{-k}, r_x] r_a^{-i} = [r_b^m r_y^{-1}, r_x] r_a^{-i} = [r_y^{-1}, r_x] r_a^{-i} = [r_y^{-1}, r_x]. \end{aligned}$$

So  $u_{xy} r_y^{-1} = r_x^{-1} r_y^{-1} r_x = r_y^{-1} [r_y^{-1}, r_x] = r_y^{-1} u_{xy}$ . Thus  $[u_{xy}, r_y] = 1$ . Similarly,  $u_{yx} = [r_x^{-1}, r_y]$ . This, together with  $[r_y, r_x^{-1}] = [r_x^{-1}, r_y]^{-1} = u_{yx}^{-1} = u_{xy}$  implies that  $[u_{xy}, r_x] = 1$ . Now,  $[r_x^i, r_y] = [r_x, r_y]^i = u_{xy}^i$  for all  $i \geq 0$ . So  $u_{xy}^p = 1$  and  $\langle r_x, r_y \rangle = p_+^{1+2}$ .  $\square$

**Proposition 2.7.** *Assume that (C1),  $\dots$ , (C4) hold in  $S$ . Then  $R' \leq Z(R)$  and  $|R'| \leq p$ .*

*Proof.* For  $x, y \in P$ , let  $U_{xy} = \langle u_{xy} \rangle$ . Let  $a, b$  be adjacent in  $\Gamma(H'(x))$  and  $ab \cap H(x) = \{c\}$ . Now  $r_b = r_a^i r_c^j$  for some  $i, j$ ,  $1 \leq i, j \leq p-1$ . Since  $[r_x, r_c] = 1$ , we have

$$u_{xb} = [r_x, r_b] = [r_x, r_a^i r_c^j] = [r_x, r_a^i] = [r_x, r_a]^i = u_{xa}^i.$$

So  $U_{xb} = U_{xa}$ . This, together with (C2), implies that  $U_{xy}$  is independent of the choice of  $y$  in  $H'(x)$ . Since  $u_{xy} = u_{yx}^{-1}$ , we have  $U_{xy} = U_{yx}$ . So, if  $x, y \in P$  with  $y \in H'(x)$ , then  $U_{xy} = U_{yx}$ . Now, by (C4),  $U_{xy}$  is independent of the edge  $\{x, y\}$  in  $\Sigma(P)$ . We denote this common subgroup by  $U$ .

We now show that  $U \leq Z(R)$ . Let  $x \in P$  and  $y \in H'(x)$ . We show that  $[u_{xy}, r_z] = 1$  for each  $z \in P$ . We may assume that  $z \in H'(x) \cup H'(y)$ . In this case it is clear from Proposition 2.6 because  $U_{xy} = U_{xz}$  if  $z \in H'(x)$ . Similarly, if  $z \in H'(y)$ .

Now, since  $R = \langle r_x : x \in P \rangle$ ,  $u_{xy} \in Z(R)$  and  $u_{xy} = 1$  if  $y \in H(x)$ , it follows that  $R' = \langle u_{xy} : x \in P, y \in H'(x) \rangle = U$  and is of order at most  $p$  (Proposition 2.6).  $\square$

**Proposition 2.8.** *Assume that (C1),  $\dots$ , (C4) hold in  $S$ . If  $R$  is nonabelian then exponent of  $R$  is  $p$  or  $4$  according as  $p$  is odd or  $p = 2$ . In particular, if  $P$  is finite then  $R$  is finite and  $\Phi(R) = R'$ .*

*Proof.* Let  $r = r_1 r_2 \cdots r_n \in R$ ,  $r_i \in R_\psi$ . We use induction on  $n$ . Let  $r = h r_n$ , where  $h = r_1 r_2 \cdots r_{n-1}$ . Since  $R' \subseteq Z(R)$ ,  $r_n^i h = h r_n^i [r_n^i, h] = h r_n^i [r_n, h]^i$ . So  $r^{i+1} = h^{i+1} r_n^{i+1} [r_n, h]^{1+2+\cdots+i}$  for all  $i \geq 0$ . Now, the result follows because by induction  $h^p = 1$  if  $p$  is odd and  $h^4 = 1$  if  $p = 2$ . Note that if  $p = 2$ , exponent of  $R$  can not be 2 as  $R$  is nonabelian.

Now, if  $P$  is finite then  $R/R'$  and so  $R$  are finite and  $\Phi(R) = R' \langle r^p : r \in R \rangle = R'$ . For  $p = 2$ , the last equality holds because  $r^2 \in R'$  for every  $r \in R$ .  $\square$

We now summarize the above results.

**Theorem 2.9.** *Let  $S = (P, L)$  be a connected partial linear space of prime order  $p$ . Suppose that for each  $x \in P$  there is associated a geometric hyperplane  $H(x)$  containing  $x$  such that (C1),  $\dots$ , (C4) hold. Let  $(R, \psi)$  be a nonabelian representation of  $S$  such that  $[\psi(x), \psi(y)] = 1$  for all  $x, y \in P$  with  $y \in H(x)$ . Then the following hold:*

- (i) *If  $x, y \in P$  with  $y \in H'(x)$ , then  $[\psi(x), \psi(y)] \neq 1$  and  $\langle \psi(x), \psi(y) \rangle = p_+^{1+2}$ ;*
- (ii)  *$|R'| = p$ ,  $R' \subseteq Z(R)$ ,  $R$  is a  $p$ -group, and exponent of  $R$  is  $p$  or  $4$  according as  $p$  is odd or  $p = 2$ .*

*Further,  $R_\psi \cap Z(R) = \{1\}$ ;  $\psi$  is faithful if  $H(x) \neq H(y)$  whenever  $x \not\sim y$ ; and  $R$  is finite with  $R' = \Phi(R)$  if  $P$  is finite.*

**Remark 2.10.** *For  $p = 2$ , Theorem 2.9(ii) is a consequence of ([12], Lemma 3.5, p. 310) where Ivanov did not assume (C3). Our proof of Proposition 2.7 is similar to that of ([13], Lemma 2.2, p. 526).*

**Corollary 2.11.** *Let  $S$  and  $(R, \psi)$  be as in Theorem 2.9. If  $P$  is finite then  $(R, \psi)$  is the cover of a representation  $(R_1, \psi_1)$  of  $S$  where  $R_1$  is extraspecial or  $p = 2$  and  $Z(R_1)$  is cyclic of order 4.*

*Proof.* If  $Z(R)$  is elementary abelian (this is the case if  $p$  is odd), write  $Z(R) = R'T$ ,  $R' \cap T = \{1\}$  for some subgroup  $T$  of  $Z(R)$ . Let  $R_1 = R/T$ . Then  $R_1$  is extra special. Define  $\psi_1$  from  $P$  to  $R_1$  by  $\psi_1(x) = \langle r_x T \rangle$ ,  $x \in P$ . Since  $r_x \notin Z(R)$ ,  $\langle r_x T \rangle$  is a subgroup of  $R_1$  of order  $p$  for each  $x \in P$ . Then  $(R_1, \psi_1)$  is a nonabelian representation of  $S$  and  $(R, \psi)$  is a cover of  $(R_1, \psi_1)$ .

If  $Z(R)$  is not elementary abelian, then  $p = 2$ . Write  $Z(R) = \langle a \rangle K$ ,  $\langle a \rangle \cap K = \{1\}$  where  $K \leq Z(R)$  and  $a$  is of order 4. Since  $r^2 \in R'$  for every  $r \in R$ , it follows that  $R' = \langle a^2 \rangle$ . Now taking  $R_1 = R/K$ , the above argument completes the proof.  $\square$

### 3. NONABELIAN REPRESENTATION GROUP OF A POLAR SPACE

If a polar space of rank  $r \geq 2$  and of order  $p$  admits a faithful abelian representation then the polar space is necessarily classical (for rank 2 case, see [17], 4.4.8, p. 76) and the representation is, up to a projective linear transformation, a standard one. The following proposition shows that a polar space of finite rank and of order  $p$  admits a nonabelian representation only if  $p$  is odd. For any representation  $(R, \psi)$  of  $S$ , Definition 1.1(ii) implies that  $[r_x, r_y] = 1$  if  $y \in x^\perp$ . By Example 2.1, all the results of the previous section hold.

**Proposition 3.1.** *Let  $S = (P, L)$  be a polar space of finite rank  $r \geq 2$  and of order three. Then every representation of  $S$  is abelian.*

*Proof.* Let  $(R, \psi)$  be a representation of  $S$ . By Lemma 1.3, there exists a chain of subspaces  $Q_0 = P \supsetneq Q_1 \supsetneq Q_2 \supsetneq \dots \supsetneq Q_{r-2}$  such that  $Q_i$  is a polar space of rank  $r - i$ . Thus  $Q_{r-2}$  is a

(2,  $t$ )-GQ. Let  $x, y \in Q_{r-2}$ ,  $x \approx y$ , and  $T$  be a (2, 1)-GQ in  $Q_{r-2}$  containing  $x$  and  $y$ . Such a  $T$  exists because each line has 3 points. Let  $\{x, y\}^\perp = \{a, b\}$  in  $T$ . For  $u \sim v$ , we define  $u * v \in P$  by  $uv = \{u, v, u * v\}$ . In  $T$ , since  $[r_b, r_y] = [r_b, r_x] = 1$  and  $r_{(a*x)*(b*y)} = r_{(a*y)*(b*x)}$ , it follows that  $r_x r_y = r_y r_x$ . Now, Corollary 2.5(i) completes the proof.  $\square$

*For the rest of this paper we assume that  $p$  is an odd prime.*

Let  $S = (P, L)$  be a polar space of finite rank  $r \geq 2$  and of order  $p$  and  $(R, \psi)$  be a nonabelian representation of  $S$ . Note that if  $r \geq 3$ , then finiteness of  $P$  and that of  $r$  are equivalent. However, if  $S$  is a GQ with  $s + 1$  points per line, then finiteness of  $P$  is not known except when  $s = 2, 3, 4$  (see [3], p.100). The rest of this section is devoted to prove that  $R$  is extraspecial if  $P$  is finite.

**Lemma 3.2.**  *$\psi$  is faithful and  $[r_x, r_y] \neq 1$  if  $x \approx y$ .*

*Proof.* This follows from Corollary 2.5(i) and (iv).  $\square$

Given a line  $l$  and two distinct points  $a$  and  $b$  on it, we write

$$\psi(l) = \{\langle r_a \rangle, \langle r_b \rangle, \langle r_a r_b \rangle, \langle r_a^2 r_b \rangle, \dots, \langle r_a^{p-1} r_b \rangle\}.$$

Let  $x, y \in P$ ,  $x \approx y$  and  $u, v \in \{x, y\}^\perp$ ,  $u \approx v$ . Then  $[r_x, r_y] \neq 1$  and  $[r_u, r_v] \neq 1$ . Let  $l_0 = xu$ ,  $l_1 = vy$ ,  $m_0 = xv$  and  $m_1 = uy$ . Consider the lines  $l_0$  and  $l_1$ . By ‘one or all’ axiom, each point of  $l_0$  is collinear with exactly one point of  $l_1$  and vice-versa. Let  $l_0 = \{x, u, x_1, x_2, \dots, x_{p-1}\}$  and  $\langle r_{x_i} \rangle = \langle r_x^i r_u \rangle$  for  $1 \leq i \leq p-1$ . Let  $x_i \sim v_i$  in  $l_1$ . Then  $l_1 = \{v, y, v_1, v_2, \dots, v_{p-1}\}$ . Replacing the generator  $r_v$  by  $r_v^j$  for some  $j$  ( $2 \leq j \leq p-1$ ), if necessary, we may assume that  $\langle r_{v_1} \rangle = \langle r_v r_y \rangle$ . So  $[r_x r_u, r_v r_y] = 1$ . Then  $[r_x^i r_u, r_v^i r_y] = 1$  for all  $i \geq 0$  because  $R' \subseteq Z(R)$ . By Lemma 3.2,  $[r_x^i r_u, r_v^j r_y] \neq 1$  if  $i \neq j$ . So  $\langle r_{v_i} \rangle = \langle r_v^i r_y \rangle$ . Let  $m_{i+1}$  be the line such that  $\psi(m_{i+1}) = \langle r_x^i r_u, r_v^i r_y \rangle$ ,  $1 \leq i \leq p-1$ .

Let  $z \in m_i \setminus (l_0 \cup l_1)$  and  $w \in m_j \setminus (l_0 \cup l_1)$  for  $i \neq j$ ,  $0 \leq i, j \leq p$ . If  $i = 0$ , then  $\langle r_z \rangle = \langle r_x^{k_1} r_u \rangle$  and if  $i > 0$  then  $\langle r_z \rangle = \langle (r_x^{i-1} r_u)^{k_1} (r_v^{i-1} r_y) \rangle$  for some  $k_1$ ,  $1 \leq k_1 \leq p-1$ . Similarly,  $\langle r_w \rangle = \langle r_x^{k_2} r_u \rangle$  or  $\langle (r_x^{j-1} r_u)^{k_2} (r_v^{j-1} r_y) \rangle$  for some  $k_2$ ,  $1 \leq k_2 \leq p-1$ , according as  $j = 0$  or  $j > 0$ . Now, from  $R' \subseteq Z(R)$ , the identity  $[r_x, r_y] = [r_v, r_u]$  (a consequence of  $[r_x r_u, r_v r_y] = 1$ ) and the fact that each point of  $m_i$  is collinear with exactly one point of  $m_j$  for  $i \neq j$  (a consequence of ‘one or all’ axiom), the following lemma is straight forward.

**Lemma 3.3.**  *$z \sim w$  if and only if  $k_1 + k_2 = p$ .*

**Proposition 3.4.** *If  $a, d \in R_\psi$  then  $ad[a, d]^{(p-1)/2} \in R_\psi$ .*

*Proof.* Let  $a, d \in R_\psi - \{1\}$ . Let  $x_1, x_2 \in P$  be such that  $\langle r_{x_1} \rangle = \langle a \rangle$  and  $\langle r_{x_2} \rangle = \langle d \rangle$ . We may assume that  $x_1 \approx x_2$ . Then  $[a, d] \neq 1$  by Lemma 3.2. We show that  $\langle ad[a, d]^{(p-1)/2} \rangle$  is the image of some element of  $P$ . Let  $y_1, y_2 \in \{x_1, x_2\}^\perp$  be such that  $y_1 \approx y_2$ ,  $\langle r_{y_1} \rangle = \langle b \rangle$  and  $\langle r_{y_2} \rangle = \langle c \rangle$ . Consider

the lines  $l_0 = x_1y_1$  and  $l_1 = x_2y_2$ . Let  $z_1 \in l_0$  be such that  $\langle r_{z_1} \rangle = \langle ab \rangle$  and let  $z_1 \sim z_2 \in l_1$ . Replacing the generator  $c$  by  $c^j$  for some  $j$ , if necessary, we may assume that  $\langle r_{z_2} \rangle = \langle cd \rangle$ . Let  $m_0 = x_1y_2$  and  $m_1 = z_1z_2$ . Let  $u \in m_0$  be such that  $\langle r_u \rangle = \langle a^{(p-1)/2}c \rangle$ . Then  $x_1 \neq u \neq y_2$ . Let  $u \sim v$  in  $m_1$ . By Lemma 3.3,  $\langle r_v \rangle = \langle (ab)^{(p+1)/2}(cd) \rangle$ . If  $y_1 \sim w$  in the line  $uv$ , then  $\langle r_w \rangle = \langle (a^{(p-1)/2}c)^k (ab)^{(p+1)/2}(cd) \rangle$  for some  $k$  ( $1 \leq k \leq p-1$ ). Now  $\left[ b, (a^{(p-1)/2}c)^k (ab)^{(p+1)/2}(cd) \right] = 1$ . So,  $[b, c]^{k+1} = 1$  and  $k+1 = p$ . The subgroup  $\langle b^{(p-1)/2} (a^{(p-1)/2}c)^{p-1} (ab)^{(p+1)/2}(cd) \rangle$  is the image of some point of  $y_1w$ . But  $b^{(p-1)/2} (a^{(p-1)/2}c)^{p-1} (ab)^{(p+1)/2}(cd) = ad[b, c]^{(p+1)/2} = ad[a, d]^{(p-1)/2}$ . In the last equality we have used  $[a, d] = [b, c]^{-1}$ , a consequence of  $[ab, cd] = 1$ . Thus,  $ad[a, d]^{(p-1)/2} \in R_\psi$ .  $\square$

**Proposition 3.5.**  $R_\psi$  is a complete set of coset representatives of  $R'$  in  $R$ .

*Proof.* Let  $r_1R' = r_2R'$  for some  $r_1, r_2 \in R_\psi$ . Since  $R' \subseteq Z(R)$ ,  $r_1$  and  $r_2$  are both trivial or are both nontrivial (Corollary 2.5(ii)). Assume that the later holds and that  $r_1 = r_2w$  for some  $w \in R'$ . Let  $x_1, x_2 \in P$  be such that  $\langle r_{x_1} \rangle = \langle r_1 \rangle$  and  $\langle r_{x_2} \rangle = \langle r_2 \rangle$ . Since  $[r_1, r_2] = 1$ , either  $x_1 = x_2$  or  $x_1 \sim x_2$  (Lemma 3.2). If  $x_1 \sim x_2$  then  $w \neq 1$  by Definition 1.1(ii) and  $\langle w \rangle$  would be the image of some point in the line  $x_1x_2$ , a contradiction to Corollary 2.5(ii). So  $x_1 = x_2$  and  $r_1 = r_2^i$  for some  $i$  ( $1 \leq i \leq p-1$ ). Then  $r_2^{i-1} = w \in R' \subseteq Z(R)$ . Now, Corollary 2.5(ii) implies that  $i = 1$  and so  $w = 1$  and  $r_1 = r_2$ .

Now, let  $sR' \in R/R'$ . Write  $s = r_1r_2 \cdots r_k$ ,  $r_i \in R_\psi$ . Let  $R' = \langle z \rangle$ . Since  $R' \subseteq Z(R)$ , there is some integer  $j$  such that  $r_1r_2 \cdots r_kz^j$  is an element, say  $r$ , of  $R_\psi$  by Proposition 3.4. Then  $sR' = rR'$ , completing the proof of the proposition.  $\square$

**Proposition 3.6.** Assume that  $P$  is finite. Then  $|R| = p(1 + (p-1)|P|)$  and  $R = p_+^{1+2m}$  for some  $m \geq 1$ .

*Proof.* Since  $|R'| = p$  (Proposition 2.7), the first assertion follows from Proposition 3.5. Also,  $R' = Z(R)$  because  $R_\psi \cap Z(R) = \{1\}$  and  $R' \subseteq Z(R)$ . Now, Proposition 2.8 completes the proof.  $\square$

**Corollary 3.7.** If  $S$  is a finite classical polar space of rank  $r \geq 2$  admitting a nonabelian representation, then  $S$  is isomorphic to  $W_{2m}(p)$  or  $Q_{2m+1}(p)$ .

*Proof.* By Proposition 3.6,  $|P| = (p^{2m} - 1)/(p - 1)$  for some  $m > 0$ . So the corollary follows from the number of points of classical polar spaces (see 1.2).  $\square$

By proposition 3.5,  $S$  admits a faithful abelian representation with representation group  $R/R'$ . Considering  $R/R'$  as a vector space over  $F_p$ , it has dimension  $2m$ . Since  $Q_{2m+1}(p)$  does not possess faithful abelian  $2m$ -dimensional representation, the only possibility is that  $S$  is isomorphic to  $W_{2m}(p)$ . We thank the referee for this remark. In the next sections, we prove this fact giving a geometrical argument involving triads of points of a generalized quadrangle.

#### 4. Rank 2 Case

Let  $S = (P, L)$  be a finite  $(s, t)$ -GQ. A *triad of points* in  $S$  is a triple  $T$  of pairwise noncollinear points. An element of  $T^\perp$  is a *center* of  $T$ . A pair of distinct points  $\{x, y\}$  in  $S$  is *regular* if  $x \sim y$  or if  $x \approx y$  and  $|\{x, y\}^{\perp\perp}| = t + 1$ . A point  $x$  is *regular* if  $\{x, y\}$  is regular for each  $y \in P \setminus \{x\}$ . The pair  $\{x, y\}$ ,  $x \approx y$ , is *antiregular* if  $|z^\perp \cap \{x, y\}^\perp| \leq 2$  for each  $z \in P \setminus \{x, y\}$ . A point  $x$  is *antiregular* if  $\{x, y\}$  is antiregular for each  $y \in P \setminus \{x\}$ . Dually, we define a triad of lines, center of a triad of lines, regularity and antiregularity of a line.

**Proposition 4.1.** *Let  $S = (P, L)$  be a  $(p, t)$ -GQ. If  $S$  admits a triad of lines with at least 3 centers then every representation of  $S$  is abelian.*

*Proof.* Let  $\{l_1, l_2, l_3\}$  be a triad of lines in  $S$  with centers  $m_1, m_2, m_3$ . Let  $\{x_{ij}\} = l_i \cap m_j$ ,  $1 \leq i, j \leq 3$ . Consider the lines  $l_1$  and  $l_2$ . Replacing  $r_{x_{11}}$  by  $r_{x_{11}}^k$  for some  $k$ , if necessary, we may assume that the point  $a$  of  $l_1$  with  $\langle r_a \rangle = \langle r_{x_{11}} r_{x_{12}} \rangle$  is collinear with the point  $b$  with  $\langle r_b \rangle = \langle r_{x_{21}} r_{x_{22}} \rangle$ . So  $[r_{x_{11}} r_{x_{12}}, r_{x_{21}} r_{x_{22}}] = 1$ . Then  $[r_{x_{11}}^i r_{x_{12}}, r_{x_{21}}^i r_{x_{22}}] = 1$  for  $0 \leq i \leq p - 1$ . Let  $\langle r_{x_{13}} \rangle = \langle r_{x_{11}}^i r_{x_{12}} \rangle$  and  $\langle r_{x_{23}} \rangle = \langle r_{x_{21}}^j r_{x_{22}} \rangle$  for some  $i, j$ ,  $1 \leq i, j \leq p - 1$ . If  $i \neq j$  then  $R$  is abelian (Corollary 2.5(i)). So assume that  $i = j$ . Let  $\langle r_{x_{31}} \rangle = \langle r_{x_{11}}^k r_{x_{21}} \rangle$  and  $\langle r_{x_{33}} \rangle = \langle (r_{x_{11}}^i r_{x_{12}})^n (r_{x_{21}}^i r_{x_{22}}) \rangle$  for some  $k, n$ ,  $1 \leq k, n \leq p - 1$ . If  $n \neq p - k$ , then  $R$  is abelian by Lemma 3.3. So, we assume that  $\langle r_{x_{33}} \rangle = \langle (r_{x_{11}}^i r_{x_{12}})^{p-k} (r_{x_{21}}^i r_{x_{22}}) \rangle$ . By a similar argument, we assume that  $\langle r_{x_{32}} \rangle = \langle r_{x_{21}}^{p-k} r_{x_{22}} \rangle$ . Now, Lemma 3.3 implies that  $R$  is abelian because  $x_{32} \sim x_{33}$  and  $p - k \neq p - (p - k)$ .  $\square$

**Corollary 4.2.** *If  $S$  admits a nonabelian representation then every line of  $S$  is antiregular and no line of  $S$  is regular.*

**Proposition 4.3.** *Let  $S = (P, L)$  be a finite  $(p, t)$ -GQ. If  $S$  admits a nonabelian representation  $(R, \psi)$ , then  $t = p$  and  $R = p_+^{1+4}$ .*

*Proof.* We have  $|P| = (p + 1)(pt + 1)$  ([17], 1.2.1, p. 2). So  $|R| = p^2(t(p^2 - 1) + p)$  (Proposition 3.6). By Corollary 4.2,  $t \geq 2$ . So,  $p^2(t(p^2 - 1) + p) \geq p^4$ . Now,  $|R| = p^{2m+1}$  for some integer  $m \geq 1$ . Thus,

$$t = p \left( p^{2(m-2)} + p^{2(m-3)} + \dots + p^2 + 1 \right).$$

Since  $t \leq p^2$  ([17], 1.2.3, p. 3),  $m = 2$ ,  $t = p$  and  $R = p_+^{1+4}$ .  $\square$

In  $Q_5(p)$  all lines are regular ([17], 3.3.1(i), p 51). So every representation of  $Q_5(p)$  is abelian. On the other hand, since  $p$  is odd,  $W_4(p)$  is not self-dual and is isomorphic to the dual of  $Q_5(p)$  ([17], 3.2.1, p. 43). No point of  $Q_5(p)$  is regular ([17], 1.5.2(i), p. 13), so no line of  $W_4(p)$  is regular. Again, all points of  $Q_5(p)$  are antiregular ([17], 3.3.1(i), p. 51), so all lines of  $W_4(p)$  are antiregular. We prove

**Proposition 4.4.** *Let  $S = (P, L)$  be a  $(p, p)$ -GQ. If  $S$  admits a nonabelian representation then  $S$  is isomorphic to  $W_4(p)$ .*

*Proof.* Since  $W_4(p)$  is characterized by the regularity of each of its point ([17], 5.2.1, p. 77), it is enough to show that if  $x, y \in P$  and  $x \approx y$  then  $\{x, y\}^{\perp\perp}$  contains  $\{a, b\}^\perp$  for distinct  $a, b \in \{x, y\}^\perp$ . Let  $(R, \psi)$  be a nonabelian representation of  $S$ . Let  $z \in \{a, b\}^\perp$  and  $w \in \{x, y\}^\perp$ . We claim that  $z \sim w$ . Write  $H = C_R(r_a) \cap C_R(r_b)$ . Then

$$|H| = \frac{|C_R(r_a)| |C_R(r_b)|}{|C_R(r_a) C_R(r_b)|} = \frac{p^4 p^4}{p^5} = p^3.$$

Let  $K = \langle r_x, r_y \rangle$ . By Proposition 2.6,  $|K| = p^3$ . So  $K = H$  because  $K \leq H$ . Then  $[r_w, r_z] = 1$  because  $[r_w, K] = 1$ . So  $z \sim w$  by Theorem 2.9(i).  $\square$

## 5. Proof of Theorems 1.5 and 1.6

**Proof of Theorem 1.5.** By Proposition 3.1,  $p$  is an odd prime. By Lemma 1.3 and Proposition 4.4,  $S$  is isomorphic to  $W_{2r}(p)$ . Proposition 3.6 implies that  $R = p_+^{1+2r}$ . This completes the proof of Theorem 1.5.

We prove Theorem 1.6 in Propositions 5.2 and 5.3. In view of Proposition 3.4, we first prove

**Proposition 5.1.** *Let  $G = p_+^{1+2r}$ . There exists a set  $T$  of coset representatives of  $Z(G)$  in  $G$  such that if  $t_1, t_2 \in T$  then  $t_1 t_2 [t_1, t_2]^{(p-1)/2} \in T$ . Further,  $T$  is unique up to conjugacy in  $G$ .*

*Proof.* Let  $Z = Z(G) = \langle z \rangle$  and  $V = G/Z$ . We consider  $V$  as a vector space over  $F_p$ . The map  $f : V \times V \rightarrow F_p$  taking  $(xZ, yZ)$  to  $i$ , where  $[x, y] = z^i$  ( $0 \leq i \leq p-1$ ), is a nondegenerate symplectic bilinear form on  $V$ . Write  $V$  as an orthogonal direct sum of  $r$  hyperbolic planes  $K_i$  ( $1 \leq i \leq r$ ) in  $V$  and let  $H_i$  be the inverse image of  $K_i$  in  $G$ . Then  $H_i$  is generated by 2 elements  $x_{i_1}$  and  $x_{i_2}$  such that  $[x_{i_1}, x_{i_2}] = z$ . Let  $A_j = \langle x_{i_j}, 1 \leq i \leq r \rangle$ ,  $j = 1, 2$ . Then  $A_j$  is an elementary abelian  $p$ -subgroup of  $G$  of order  $p^r$ ,  $A_j \cap Z = \{1\}$  and  $A_1 Z \cap A_2 Z = Z$ . Set

$$T = \left\{ xy [x, y]^{\frac{p-1}{2}} : x \in A_1, y \in A_2 \right\}.$$

We show that  $T$  has the required property. Let  $\alpha = xy [x, y]^{\frac{p-1}{2}}$ ,  $\beta = uv [u, v]^{\frac{p-1}{2}}$  be elements of  $T$  where  $x, u \in A_1$  and  $y, v \in A_2$ . If  $\alpha Z = \beta Z$ , then  $u^{-1}xZ = y^{-1}vZ$  and is equal to  $Z$  because  $A_1 Z \cap A_2 Z = Z$ . So  $x = u$  and  $y = v$  because  $A_j \cap Z = \{1\}$ . Thus  $\alpha Z = \beta Z$  if and only if  $x = u, y = v$ . So,  $|T| = p^{2r}$  and  $T$  is a complete set of coset representatives. Since  $G' = Z$ , a routine calculation shows that  $\alpha\beta [\alpha, \beta]^{(p-1)/2} = (xu)(yv)[xu, yv]^{(p-1)/2} \in T$ . Thus,  $T$  has the stated property.

Now we prove the uniqueness part. In fact, we show that the group of inner automorphisms of  $G$  acts regularly on the set  $\mathcal{X}$  of all sets of coset representatives of  $Z$  in  $G$ , each of which is closed under the binary operation  $(t_1, t_2) \mapsto t_1 t_2 [t_1, t_2]^{(p-1)/2}$ .

Fix an ordered basis  $\{v_1 Z, \dots, v_{2r} Z\}$  for  $V$ . Each  $T \in \mathcal{X}$  is determined by the sequence  $(x_1, \dots, x_{2r})$ , where  $T \cap v_i Z = \{x_i\}$ . In fact, if  $aZ = x_{i_1}^{j_1} \dots x_{i_n}^{j_n} Z \in V$ , where  $i_1 < \dots < i_n$  and  $1 \leq j_k \leq p-1$ , then  $aZ \cap T = \{x_{i_1}^{j_1} \dots x_{i_n}^{j_n} z^m\}$ , where

$$z^m = [x_{i_1}^{j_1}, x_{i_2}^{j_2}]^{(p-1)/2} [x_{i_1}^{j_1} x_{i_2}^{j_2}, x_{i_3}^{j_3}]^{(p-1)/2} \dots [x_{i_1}^{j_1} \dots x_{i_{n-1}}^{j_{n-1}}, x_{i_n}^{j_n}]^{(p-1)/2}.$$

Thus,  $|\mathcal{X}| \leq p^{2r}$ . Further, for  $T \in \mathcal{X}$  and  $g \in G$ ,  $g^{-1}Tg = T$  implies  $g \in Z$ . To see this, let  $t \in T$  and  $g^{-1}tg = t' \in T$ . Then,  $tZ = g^{-1}tgZ = t'Z$ . Since  $T$  contains exactly one element from each coset, it follows that  $t = t'$  and  $g \in C_G(t)$ . Thus,  $g \in C_G(T) = Z$ . Since  $|G : Z| = p^{2r}$ ,  $|\mathcal{X}| = p^{2r}$  and  $G$  acts transitively on  $\mathcal{X}$ .  $\square$

**Proposition 5.2.**  $W_{2r}(p)$ ,  $r \geq 2$ , admits a nonabelian representation and the representation group is  $p_+^{1+2r}$ .

*Proof.* Let  $G = p_+^{1+2r}$  and  $T$  be as in Proposition 5.1. Consider the partial linear space  $S = (P, L)$ , where  $P = \{\langle x \rangle : 1 \neq x \in T\}$  and a line is of the form  $\{\langle x \rangle, \langle y \rangle, \langle xy \rangle, \dots, \langle x^{p-1}y \rangle\}$  for distinct  $\langle x \rangle, \langle y \rangle$  in  $P$  with  $[x, y] = 1$ . Note that  $x^i y \in T$  for each  $i$  and  $|P| = (p^{2r} - 1)/(p - 1)$ . We show that  $S$  is a polar space of rank  $r$ .

Since  $T \cap Z(G) = \{1\}$ ,  $S$  is nondegenerate. Let  $\langle x \rangle \in P$ ,  $l \in L$  and  $\langle x \rangle \notin l$ . Then,  $\langle x \rangle$  is collinear with one or all points of  $l$  because  $C_G(x)$  intersects nontrivially with the subgroup  $H$  of  $G$  generated by the points of  $l$ . Note that  $H$  is a subgroup of order  $p^2$  and disjoint from  $Z(G)$ . Rank of  $S$  is  $r$  because singular subspaces in  $S$  correspond to elementary abelian subgroups of  $G$  which intersect  $Z(G)$  trivially and  $p^r$  is the maximum of the orders of such subgroups of  $G$ . Thus  $S$  is a polar space of rank  $r$ .

Clearly  $G$  is a representation group of  $S$ . So,  $S$  is isomorphic to  $W_{2r}(p)$  (Theorem 1.5(iii)).  $\square$

**Proposition 5.3.** Any two representations of  $W_{2r}(p)$ ,  $r \geq 2$ , are equivalent.

*Proof.* Let  $(R_1, \psi_1)$  and  $(R_2, \psi_2)$  be two representations of  $W_{2r}(p)$ . By Theorem 1.5(ii), we may assume that  $R_1 = R_2 = R$ . By Proposition 3.5, each  $R_{\psi_i}$  is a set of coset representatives of  $Z(R)$  in  $R$ . Let  $\varphi \in \text{Aut}(R)$  be such that  $\varphi(R_{\psi_1}) = R_{\psi_2}$  (Proposition 5.1). Define  $\beta : P \rightarrow P$  by  $\beta = \psi_2^{-1} \varphi \psi_1$ . Now, Lemma 3.2 implies that  $\beta$  is an automorphism of  $W_{2r}(p)$ . Now,  $(R, \psi_1)$  and  $(R, \psi_2)$  are equivalent with respect to  $\varphi$  and  $\beta$ .  $\square$

## REFERENCES

- [1] A. E. Brouwer and H. A. Wilbrink, The structure of near polygons with quads, *Geom. Dedicata* **14** (1983), no. 2, 145–176.
- [2] F. Buekenhout and E. Shult, On the foundations of polar geometry, *Geom. Dedicata* **3** (1974), 155–170.

- [3] P. J. Cameron, “Projective and polar spaces”, available from <http://www.maths.qmul.ac.uk/~pjc/pps/>
- [4] A. M. Cohen and B. N. Cooperstein, A characterization of some geometries of Lie type, *Geom. Dedicata* **15** (1983), 73–105.
- [5] B. N. Cooperstein, Geometry of long root subgroups in groups of Lie type, Proc. Sympos. Pure Math., 37, 243–248, Amer. Math. Soc., Providence, R.I., 1980.
- [6] B. N. Cooperstein, The geometry of root subgroups in exceptional groups *I*, *Geom. Dedicata* **8** (1979), 317–381.
- [7] B. N. Cooperstein, The geometry of root subgroups in exceptional groups *II*, *Geom. Dedicata* **15** (1983), 1–45.
- [8] K. Doerk and T. Hawkes, “Finite soluble groups,” de Gruyter Expositions in Mathematics 4, Walter de Gruyter & Co., Berlin, 1992.
- [9] D. Frohardt and P. M. Johnson, Geometric hyperplanes in generalized hexagons of order  $(2, 2)$ , *Comm. Algebra* **22** (1994), no. 3, 773–797.
- [10] R. Gramlich, On graphs, geometries, and groups of Lie type, PhD Thesis, Technische Universiteit, Eindhoven, 2002.
- [11] R. L. Griess, Jr., The Monster and its nonassociative algebra, in “Finite groups—coming of age” (Montreal, Que., 1982), 121–157, Contemp. Math., 45, Amer. Math. Soc., Providence, RI, 1985.
- [12] A. A. Ivanov, Non-abelian representations of geometries, in “Groups and combinatorics—in memory of Michio Suzuki”, 301–314, Adv. Stud. Pure Math., 32, Math. Soc. Japan, Tokyo, 2001.
- [13] A. A. Ivanov, D. V. Pasechnik and S. V. Shpectorov, Non-abelian representations of some sporadic geometries, *J. Algebra* **181** (1996), no. 2, 523–557.
- [14] P. M. Johnson, Polar spaces of arbitrary rank, *Geom. Dedicata* **35** (1990), no. 1-3, 229–250.
- [15] W. M. Kantor, Generalized quadrangles having a prime parameter, *Israel J. Math.* **23** (1976), no. 1, 8–18.
- [16] S. E. Payne, A census of finite generalized quadrangles, in “Finite geometries, buildings, and related topics” (Pingree Park, CO, 1988), 29–36, Oxford Sci. Publ., Oxford Univ. Press, New York, 1990.
- [17] S. E. Payne and J. A. Thas, “Finite generalized quadrangles,” Research Notes in Mathematics, 110, Pitman (Advanced Publishing Program), Boston, MA, 1984.
- [18] B. K. Sahoo and N. S. N. Sastry, On the order of a non-abelian representation group of a slim dense near hexagon, communicated.
- [19] J. Sarli, The geometry of root subgroups in Ree groups of type  ${}^2F_4$ , *Geom. Dedicata* **26** (1988), 1–28.
- [20] J. A. Thas, Generalized polygons, in “Handbook of incidence geometry,” 383–431, North-Holland, Amsterdam, 1995.
- [21] J. A. Thas, Projective geometry over a finite field, in “Handbook of incidence geometry,” 295–347, North-Holland, Amsterdam, 1995.
- [22] J. Tits, “Buildings of spherical type and finite BN-pairs,” Lecture Notes in Mathematics, Vol. 386. Springer-Verlag, Berlin-New York, 1974.